



stability-indicating assay method for determination of Tinidazole in active substance with Spectroscopic approach

Mostafa Khajeh Dehaghani

Student of pharmaceutical supervision at Tehran University of Medical Sciences

Abstract

A new, simple, rapid, accurate, sensitive and cost-effective spectrophotometric stability indicating assay method was developed for the quantification of Tinidazole in bulk and validated according to ICH guidelines using different parameters. In the present study an attempted has been made to develop an effective method which will surpass the disadvantages associate with other reported methods like tedious in use, less sensitive and costly etc. The assay method was based on the forced degradation study of Tinidazole under different ICH recommended stress conditions like hydrolytic, oxidation, thermal, and photolysis. Drug shown maximum absorption at 322nm and a regression coefficient value (R^2) of 0.999 indicates linearity followed within the concentration range of 5 μ g/ml to 50 μ g/ml. During forced degradation study, Tinidazole was significantly degraded in alkaline conditions whereas mild degradation occurs in acidic, neutral and oxidative stress. The drug was stable to dry heat and photolytic degradation conditions. Statistical validation result of the developed method indicates that it was a simple, precise, reproducible, selective, specific and accurate method for analysis of Tinidazole and could be successfully adopted to estimate the drug in bulk and formulations.

Keywords: Tinidazole; Stability indicating assay method; UV spectroscopy; Validation; Forced degradation study; impurity

Introduction

Tinidazole (TIN) chemically known as 1-(2-ethylsulfonyl-ethyl)-2-methyl-5-nitroimidazole having molecular formula and molecular mass $C_8H_{13}N_3O_4S$ and 247.27 g/mol respectively (Fig.1)[1]. Tinidazole is an antibiotic that is used to treat certain infections caused by bacteria, such as infection of the intestines or vagina. Tinidazole is also used to treat certain sexually transmitted infections. Tinidazole is used to treat giardiasis or amebiasis in adults and children at least 3 years old. Tinidazole may also be used for purposes other than those listed in this medication guide. . Recently there is an increased need for development of a stability-indicating assay method (SIAM), using the conditions mentioned in the ICH guidelines[8] . A stability method is a validated analytical technique used to measure the concentration of active constituents accurately in the presence of its degradation products, excipients, and other impurities. The International

Conference on Harmonization (ICH) guideline entitled “Stability testing of new drug substances and products” requires that stress testing be carried out to elucidate the inherent stability characteristics of the active substance by the influence of temperature, humidity, light, oxidizing agent as well as susceptibility over a wide range of pH values[9] [10] Generally, chromatographic methods were established to monitor the changes in concentration of drug at different stressed conditions[11] [12]. Spectrophotometric stability method may have advantages over chromatographic techniques in terms of simplicity, economic and time consumption. In degradation study, the intensity of stress parameters depends on physicochemical properties of drug substance and nature of drug or product[13] [14]. In the

Pharmaceutical industry time and expenses plays an important role hence simple, cost effective and more sensitive methods are highly acceptable for routine usage of study. The present study was aimed to develop and validate a UV- spectroscopic stability indicating assay method for Tinidazole in distilled water at different stressed conditions [15]. The objective of the study is developing a suitable, efficient and economic analytical method to estimate drug in presence of its degradation product and impurities.





Figure 1 Chemical structure of Tinidazole

1. Material and methods

Chemicals and Reagents Pharmaceutical-grade API of Tinidazole was procured as a gift sample from FARABI Pharmaceutical and was used without further purification. Distilled water used as solvent during the study was prepared in the chemistry lab CPS, Puri and all other chemicals and reagents used were of analytical grade. 2.2. Instruments For developing the stability-indicating assay method of TIN, a PC controlled double beam UV-spectrophotometer (Systonic) having spectral bandwidth 3nm and of wavelength accuracy $\pm 1\text{nm}$ with 1 cm quartz cell was used for spectral estimation and absorbance measurements of sample solutions. The work was carried out in an air-conditioned room maintained at a temperature $25 \pm 2^\circ\text{C}$. A precision mantel heater (Biotech, Mumbai) with temperature regulator equipped with a reflux condenser was used for degradation study in acid, alkali and



neutral conditions. A dry-air oven was used to study the effect of dry heat. The photolytic study was carried out by exposing the drug to direct sunlight. 2.3. Preparation of standard stock solution and estimation of $\lambda_{\max}$



On the basis of solubility of TIN, distilled water was selected as solvent to carry out the whole study. A standard stock solution of 1000 $\mu\text{g/ml}$ of TIN was prepared by dissolving 100 mg of pure drug in a 100ml volumetric flask by adjusting the final volume with distilled water. Working standard solution of 100 $\mu\text{g/ml}$ of TIN was further prepared by diluting 10ml of the above solution to 100ml with distilled water. Then further dilutions of 5- 50 $\mu\text{g/ml}$ were made by diluting the required aliquot of 100 $\mu\text{g/ml}$ solution with distilled water. The solution with a concentration of 30 $\mu\text{g/ml}$ was scanned over the range of 200-450nm against blank solvent. The drug spectrum was showing maximum absorbance ($\lambda_{\max}$) at 322nm which is considered as the analytical wavelength for further study (Fig. 02).

1.4. Validation of Method

- 1.4.1. Linearity Linearity study of the drug was performed for the dilutions of concentration 5-50 $\mu\text{g/ml}$ which were scanned at 322nm and absorbance were recorded. The calibration curve was plotted between absorbance and concentration of standard solutions and the regression equation was determined. The experiment was carried out in five replicates. 2.4.2. Accuracy The accuracy of the developed method was confirmed by performing recovery study. The recovery study was carried out by standard addition method in which pre-analyzed sample of 20 $\mu\text{g/ml}$ were spiking with extra standard drug of TIN in 80,100 and 120% and absorbance recorded at 322nm. The percent of recovery of the added drug was calculated. The study is carried out in

triplicate. 2.4.3. Precision The precision of the method was established by repeatability and intermediate precision. Three solutions of concentration 10, 20 and 30 $\mu\text{g/ml}$ of TIN were prepared to perform the precision study. Repeatability Repeatability (intra-day) was assessed by analyzing the working solution of TIN, analyzed in triplicates for three different concentrations (10, 20 and 30 $\mu\text{g/ml}$) of three times a day at an interval of 1 hr. The % relative standard deviation (RSD) was calculated for absorbance obtained for these solutions. Intermediate precision Intermediate precision or inter-day precision was established by analyzing the working solution of three different concentrations (10, 20 and 30 $\mu\text{g/ml}$) of TIN for three different days. The results was reported in terms of % RSD. 2.4.4. Detection and Quantitation limit As per ICH guideline three approaches are to determine the detection limit (DL) and quantitation limit (QL), these includes visual evaluation, signal to noise ratio and the use of standard deviation of the response and the slope of the calibration curve. DL is the lowest amount of analyte in a sample, which can be detected and QL is the lowest amount of analyte which can be quantitatively determined by the developed method. The LOD and LOQ were separately determined based on the third approach and calculated using eq. 1 and eq.2, where, σ is the standard deviation of the intercept of the calibration plot and S is the slope of the calibration curve.

1.5. Forced Degradation

Studies 18-20 As per the International Conference on Harmonization (ICH) guideline, stability testing of a drug substances and products requires that stress testing to be performed for revealing of the inherent stability characteristics²⁰. In order to check the stability of TIN in different ICH prescribed forced degradable conditions like hydrolytic (acid, base and neutral), thermal, photolytic and oxidative, drug at a concentration of 1 mg ml⁻¹ was used for all degradation studies.

2.5.1. Acidic degradation study For acidic degradation study, 10 ml of 1mg ml⁻¹ standard stock solution of the TIN was added to 10 ml of 1.0N HCl and refluxed for 06hrs at 60 OC. From this 0.4ml of aliquot was taken at regular interval of time in separate 10ml volumetric

flask and adjusted the volume with distilled water after neutralization with 1.0N NaOH to form a solution having concentration 20 $\mu\text{g/ml}$ and absorbance was recorded at 322nm against blank solvent. Finally the absorbance of the sample were compared with the absorbance of standard solution and percentage of degradation was calculated.

2.5.2. Alkaline degradation Study 10 ml of 1mg ml⁻¹ standard stock solution of the TIN was added to equal volume of 1.0N NaOH and refluxed for 06hrs at 60 OC. From this 0.4ml of aliquot was taken at regular interval of time in a 10ml volumetric flask and adjusted the volume by distilled water after neutralization with 1.0N HCl to form a solution containing 20 $\mu\text{g/ml}$



and absorbance was recorded at 322nm against blank solvent. The percentage of degradation was calculated by comparing the absorbance of the sample with the absorbance of standard solution.

2.5.3. Neutral Hydrolysis In separate volumetric flask 20 ml of 1 mg ml⁻¹ standard stock solution of the TIN was refluxed for 06hrs at 60 °C. From this 0.2ml of aliquot was taken regularly in separate 10ml volumetric flask which was then adjusted the volume with distilled water to form a solution containing 20µg/ml and absorbance was recorded at 322nm. Absorbance of sample and standard solutions were compared to determine the percentage of decomposition.

2.5.4. Oxidation induced degradation To perform oxidative degradation study 10 ml of 1mg ml⁻¹ standard stock solution of the TIN is mixed with 10 ml of 3% H₂O₂. The solution was kept at room temperature for a period for 24hrs. A volume of 0.4 ml of aliquot was taken after 24hrs in a separate 10ml volumetric flask and adjusted the volume with distilled water to form a solution of 20 µg/ml and at 322nm absorbance was recorded and percentage of degradation was calculated.

2.5.5. Thermal Degradation A sample power of accurately weighed 100 mg of TIN was exposed to 80 °C for 48hrs in a controlled-temperature hot air oven. Then required amount was dissolved distilled water and a solution of 20 µg/ml was prepared and the absorbance was recorded at 322nm and degradation percent was calculated.

2.5.6. Photo degradation The pure drug sample accurately weighed 100 mg was exposed to sun light for 6hrs. Further suitably diluted to 20 µg/ml by dissolving required amount of drug in distilled water and absorbance was recorded. Finally percent of degradation was estimated by comparing absorbance of sample and standard solutions.

3. Results and discussion

3.1. UV Spectroscopic method development In the present study, a stability-indicating assay method was developed and validated as per ICH guidelines for study of degradation behaviour of TIN in bulk and marketed formulations using distilled water as solvent. The wavelength 322nm was found as λ_{max} and taken as the analytical wavelength for estimation of the drug. This developed method is found to be simple and economical as compared to the previously reported HPLC methods as it required less sample preparation and solvent consumption.

3.2. Validation of the developed method


3.2.1. Linearity and range Linearity was evaluated by analysis of working standard solutions of TIN for concentration 5-50 µg/ml. The absorbance of these solutions was recorded at 322 nm and a plot was made between the absorbance and concentration of standard solutions as shown in Figure 3. The results of the regression analysis are summarized in (Table 1). The result of the study showed that the drug obeyed Beer-Lambert's law in the above concentration range with a correlation coefficient of 0.999 which



indicates that, the method is capable enough with good sensitivity. The mean absorbance was found in between 0.036-0.957.

Table 1 Regression analysis of TIN for developed method

S.NO	Parameters	Results
1	Absorption maxima $\lambda_{\max}$	322 nm
2	Beer's-Lambert's range	5-50 $\mu\text{g/ml}$
3	Regression equation	$y = 0.057x + 0.0101$
4	Correlation coefficient	0.999
5	Slope	0.057
6	Intercept	0.0101



3.2.2. Accuracy Study

Accuracy study was performed to check the sensitivity of the developed method for the estimation of TIN. The recovery study was investigated by analysing three concentrations of standard drug solution previously analysed using the standard addition technique. The standard addition technique was carried out by adding 80,100 and 120 % of the TIN concentration in the sample. The % recoveries of the three concentrations were found to be 100.24 -100.88 %. The results of % recovery and % RSD are given in Table 2. The accuracy study results indicates that, the recoveries were well within the limits and the developed method is capable enough of showing good accuracy with reproducibility for analysis of TIN. **Table 2 Result of Recovery Study**



Initial Concentration (µg/ml)	Level of spiking %	Concentration of spiked sample (µg/ml)	Absorbance ± S.D	% Recovery (n=3)	% RSD (n=3)
20	80	16	0.694 ± 0.254	100.24	0.174
20	100	20	0.819 ± 0.145	100.57	0.112
20	120	24	0.947 ± 0.351	100.88	0.092

3.2.3. Precision Study

The results of the repeatability and intermediate precision study are given in Table 3. The developed method was found to be precise as the % RSD values for repeatability and intermediate precision studies were < 2.0 indicating the proposed method is repeatable when executed in different day also.

Table 3 Result of Precision Study

Repeatability Study			Intermediate Precision study		
Sample concentration (µg/ml)	Mean Absorbance ± S.D	% RSD	Sample concentration (µg/ml)	Mean Absorbance ± S.D	% RSD
10	0.126 ± 0.012	0.521	10	0.158 ± 0.063	0.089
20	0.281 ± 0.054	0.341	20	0.265 ± 0.154	0.184
30	0.577 ± 0.023	0.254	30	0.527 ± 0.897	0.279

3.2.4. Detection and Quantitation limits

The detection limit (DL) and quantitation limit (QL) were determined as per the ICH guidelines and were found to be 0.62 and 1.85 µg/ml for TIN respectively (Table. 4). T

able 4 Result of LOD and LOQ


3.3. Result of forced Degradation Study



Table 5 Degradation studies of TIN at different induced forced stress conditions at 322nm

S. No	ICH prescribed different stressed conditions	Concentration ($\mu\text{g/ml}$)	Absorbance	Percentage of degradation
1	Acidic condition	20	0.828	15.3
2	Alkali condition	20	0.642	58.6
3	Neutral condition	20	0.879	12.3
4	Oxidation	20	0.764	24.6
5	Thermal	20	0.952	5.8
6	Photolysis	20	0.924	5.6


The induced forced degradation study of TIN was performed by exposing standard drug of TIN to different stress conditions under ICH guidelines. Result of the study indicates that TIN was undergoes significant degradation in alkali hydrolytic conditions, whereas mild degradation under acidic, neutral and oxidation conditions but stable to thermal and photolytic conditions. The results of forced degradation study summarized in (Table 5). 3.3.1. Hydrolytic studies Acid induced degradation For the acidic hydrolytic stress degradation study, TIN dissolved in 1.0 N HCl and refluxed for 06hrs at 60 OC. After that a working solution of 20 $\mu\text{g/ml}$ was prepared by further diluting with distilled water and absorbance was recorded at 322nm. The percentage of degradation was determined by comparing the absorbance of sample solution with standard solution. In strong acidic hydrolytic condition the drug was found to be moderately degraded by 25.3% (Fig.4).




Alkali induced degradation In stress degradation study under alkali conditions, TIN exposed to 1.0 N NaOH solution and refluxed for 6hrs at 60 OC. After that the solution was neutralized by mixing with equivalent volume 1.0N HCl solution and working solution 20 $\mu\text{g/ml}$




was prepared by dilute with distilled water and absorbance was determined at 322nm. The drug was found to be degraded to a significant percentage of 58.6 % (Fig. 5).




In neutral conditions the prepared drug solution was refluxed for 06hrs at 60 OC. The percentage of degradation was found to be 15.3% (Fig. 6).



3.3.2. Oxidation induced degradation In oxidative stressed condition the drug solution is mixed to a 3% H₂O₂ solution and kept at room temperature for 24hrs. A 20 µg/ml solution was prepared from the above solution by dilute with distilled water and absorbance was taken at 322nm. The result of the study indicates that the drug was degraded by 24.6% in oxidative degradable conditions (Fig.7).




3.3.3. Thermal stress study In stress degradation study under thermal condition pure sample powder of TIN exposed at 80 OC for 48hrs. There was no notable degradation was found for thermal condition about 5.8% of standard drug was found to be degraded under this stress condition (**Fig. 8**).



3.3.4.

Photolytic degradation Photo degradation study of the drug was carried out in dry form where the drug was directly exposed to the sunlight for 6hrs on a hot sunny day. After that required amount was diluted with distilled water to make a working solution 20 µg/ml and absorbance recorded at 322nm. The percentage of degradation was estimate by comparing absorbance of sample solution with standard solution. The drug was found to be stable to the exposed degradation condition and about 5.6% degradation of drug was observed for thermal degradation study (Fig. 9).



2. Conclusion

Changes in drug stability can risk patient safety by the formation of toxic degradation product(s). The stability of a drug product or a drug substance is a critical parameter which may affect purity, potency and safety. Therefore, it is essential to know the purity profile and degradation behavior of a drug substance under various degradable conditions. The proposed method presents a simple, rapid, economic and useful stability indicating assay method for the determination of Tinidazole in bulk and formulation. Majority of the reported methods employ organic solvents as diluents and suffers two disadvantages of narrow linear range and poor sensitivity. In this method distilled water is taken as diluent which makes this method inexpensive and wide linearity range with more sensitivity for drug estimation. This developed spectroscopic method was further validated as per ICH guideline. The values of % RSD for recovery and precision study are less than 2 indicates this method has accuracy, sensitivity and precision for estimation of Tinidazole. LOD and LOQ were found to be within limits. Hence this method can be easily and conveniently adopted for routine stability indicating assay of Tinidazole in different dosage formulation before going to HPLC or HPTLC methods to save the time.

References

- [1] <https://www.drugbank.ca/drugs/db00916>.
- [2] Shueb S. A., Vani R. Method Development and Validation for Simultaneous Estimation of Metronidazole, Tetracycline, Bismuth Subcitrate in its bulk and Pharmaceutical Dosage form by RP –HPLC. *World Journal of Pharmaceutical and Life Sciences*, 2019; 5(11): 190-196.
- [3] Mishra A.K., Kumar A., Mishra A., Mishra H.V. Development of ultraviolet spectroscopic method for the estimation of Metronidazole from pharmaceutical formulation. *J Nat Sci Biol Med*, 2014; 5(2):261-264.
- [4] Mahmoud M., Elkhoudary and others. Development and Optimization of HPLC Analysis of Metronidazole, Diloxanide, Spiramycin and Cliquinol in Pharmaceutical Dosage Forms Using Experimental Design, *Journal of Chromatographic Science*, 2016; 54(10): 1701–1712.
- [5] Akarim E.I., Ibrahim K.E., Adam, M.E. Studies on the Photochemical Decomposition of Metronidazole. *International Journal Pharmaceutics*, 1991; 6: 261-264.
- [6] Godfrey R., Edwards R. A Chromatographic and Spectroscopic Study of Photodegraded Metronidazole in Aqueous Solution. *Journal of Pharmaceutical Science*, 1991; 80: 212-218.



- [7] Habib M. J., Asker A.F. Complex Formation between Metronidazole and Sodium Urate: Effect of Photodegradation of Metronidazole. *Pharmaceutical Research*, 1989; 6: 58-61.
- [8] Marciniak B., Bugaj A., Kedziora W. Kinetic Studies of the Photodegradation of Nitroimidazole Derivatives in the solid state. *Pharmazie*, 1997; 52: 220-223.
- [9] Kendall A.T., Starck E., Sugden J.K. Effect of Hydroxyl Radicals on the stability of Metronidazole in Buffer Solution at pH 9.2, *International Journal of Pharmaceutics*, 1989; 57: 217-211.
- [10] ICH (2003). Stability testing of new drug substances and products, In: *Proceeding of the International Conference on Harmonisation*, ICH, Geneva.
- [11] Bakshi M., Singh S. Development of validated stability-indicating assay methods-critical review. *Journal of Pharmaceutical and Biomedical Analysis*. 2002; 28 (6):1011-1040.
- [12] Panigrahi D., Mishra G. P., Sharma R. Study of stressed degradation behaviour of drotaverine and development of a validated stability indicating HPLC assay method. *Der Pharma Chemica*. 2012; 4(3):847-853.
- [13] Sahoo P.K., Sahoo P., Mohapatra J., Panigrahi D., Patra A.K., Mishra A. Development and Validation of Stability Indicating Assay Method of Doxycycline Hyclate by using UV-Spectrophotometer. *Journal of Drug Delivery & Therapeutics*. 2023; 13(6):89-94.
- [14] Chavhan V., Ghante M. Stability Indicating UV Spectrophotometric method development and validation of Simvastatin in bulk and tablet dosage form. *J App Pharm*. 2014; 6(2): 235-246.
- [15] Ambhore J.P., Adhao V.S., Cheke R.S., Popat R.R., Gandhi S.J. Futuristic review on progress in force degradation studies and stability indicating assay method for some antiviral drugs. *GSC Biological and Pharmaceutical Sciences*. 2021; 16(01): 133-149.
- [16] Kardile K., Damle M.C. Stability Indicating UV Spectrophotometric Method for Determination of Dronedaron Hydrochloride. *Journal of Pharmaceutical Sciences and Drug Research*. 2015; 7(1): 116-119.
- [17] Ghosh M., Mondal S., Chakraborty S., Ghosh N. A Stability Indicating Method was Developed and Validation for the Estimation of Carbamazepine in Bulk and Tablet Dosage form by UV-Spectroscopic Techniques. *Journal of Drug Delivery & Therapeutics*. 2023; 13(3):85-104.
- [18] Kaur I., Wakode S., Singh H. P. Development and Validation of Stability indicating UV Spectroscopic Method for Determination of Canagliflozin in Bulk and Pharmaceutical Dosage Form. *Pharm Methods*. 2016; 7(1): 63-69.
- [19] Gandhi S.V., Patil G.R. Development and validation of Stability Indicating UV spectroscopic method for estimation of Dapsone. *International J. of Pharmacy and Biological Sciences*. 2019; 9(2): 406-412.
- [20] Chakraborty S., Sharmin S., Rony S.R., Ahmad S.A.I., Sohrab H. Stability indicating UV/Vis spectrophotometric method for Diazepam, Development and validation. *Indian J. Pharm Sci*. 2018; 80(2): 366-373.